



Ritter-Hydrolysis Cascade Strategy for the Synthesis of Substituted gamma-Lactams

Wu, Peng; Nielsen, Thomas Eiland; Clausen, Mads Hartvig

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Wu, P., Nielsen, T. E., & Clausen, M. H. (2015). *Ritter-Hydrolysis Cascade Strategy for the Synthesis of Substituted gamma-Lactams*. Abstract from 98th Canadian Chemistry Conference and Exhibition, Ottawa, Canada. <http://abstracts.csc2015.ca/00000084.htm>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Canadian Society for Chemistry | **For Our Future**

98th Canadian Chemistry Conference and Exhibition

OTTAWA, ON
JUNE 13-17, 2015

Ritter-Hydrolysis Cascade Strategy for the Synthesis of Substituted gamma-Lactams **P Wu^{\$}**
<pengwua@gmail.com>, **TE Nielsen** and **MH Clausen^{*}**, Technical University of Denmark, Denmark.

N-Acyliminium ions are reactive synthetic species that have been used in several organic reactions, such as Pictet-Spengler, Mannich and Petasis reactions. In our continued efforts to explore the synthetic utility of *N*-acyliminium ions, we report an efficient Ritter-hydrolysis cascade strategy for the synthesis of novel substituted γ -lactams. In-situ generated *N*-substituted acyliminium ions, masked as dihydroxy- γ -lactams obtained using a reductive-cyclization method, were reacted with commercially available aryl nitriles to give bicyclic unsaturated molecules. A following acidic hydrolysis reaction afforded target substituted γ -lactams with an aryl amino moiety, which is present in a wide range of natural products, biologically active molecules, and more than 300 approved drugs. A combined Ritter-hydrolysis cascade reaction yielded the target compounds in good to excellent yield. High *cis*-diastereoselectivity was observed. This reaction has shown to be tolerable to a broad selection of different *N*-substituted acyliminium ions and aryl nitriles bearing substituents of different electronic properties on the aromatic ring. This method of using *N*-acyliminium ions in Ritter-hydrolysis reactions serves as an important supplement to the current reservoir of synthetic applications of *N*-acyliminium ions.

Platinum & Gold Sponsors



TEN YEARS
OF DISCOVERY

